rine Turtle Conparison Grant



	Description	Habitat	Food	Threats
Australian Flatback (endangered)	 90 kg 1 metre long Shell is oval/round and very flat All flippers have one claw 	Waters of northern and northwest Australia and New Guinea's Gulf of Papua Nest only in Queensland	Sea cucumbers, molluscs, jellyfish, prawns, seaweed	 People harvesting eggs and shells By-catch of other fishing Pollution Climate change
Green Sea Turtle (endangered)	 230 kg 1.5 metres long Greenish-coloured fat and cartilage All flippers have one claw 	Found in all temperate and tropical seas around the world	Only sea turtle that is strictly herbivorous as an adult Sea grass and algae	Commercial harvesting for eggs, meat, skin and shell By-catch of other fishing Pollution Climate change
Hawksbill (critically endangered)	 40-60 kg 1 metre long Narrow head and hawk-like beak All flippers have two claws 	Tropical reefs and lagoons in the Atlantic, Pacific and Indian Oceans	Sponges, anemones, squid and shrimp	Harvesting for their prized shell, often called 'tortoise-shell' By-catch of other fishing Pollution Climate change
Kemp's Ridley (critically endangered)	40 kg70cm longBroad parrot-like beakHatchlings are jet black	Gulf of Mexico	Crabs, clams, mussels, shrimp, fish, sea urchins, squid, jellyfish	 Harvesting of eggs By-catch of other fishing Climate change
Leatherback (endangered)	600 kg 2 metres long Only sea turtle without a hard shell All flippers without claws Only reptile that can control its internal body temperature slightly	Found world-wide in open ocean as far north as Alaska and as far south as southern tip of Africa	Jellyfish	 Pollution (particularly plastic bags) Climate change By-catch of other fishing
Loggerhead (critically endangered)	 180 kg 1 metre long Very large head, with a powerful jaw Heart-shaped shell Front flippers have two claws, and rear flippers have two or three claws 	Found in temperate and subtropical coastal waters worldwide	Shellfish — crabs, clams and mussels	Loss of nesting habitat due to coastal development Coastal lighting that disorientates hatchlings By-catch of other fishing Climate change
Olive Ridley (critically endangered)	 35-40 kg 1 metre long All flippers have one or two claws shell is bony, with large scales (scutes) 	Tropical and subtropical waters of the Pacific, Indian and Atlantic oceans SCHOLAST ASCHOLAST	Shrimp, crabs, molluscs, tunicates and fish	 People harvesting eggs and shells Loss of nesting habitat due to coastal development By-catch of other fishing Pollution Climate change